

Can all breaks in GRB afterglows be explained by jet effects?

D.M. Wei^{1,2} and T. Lu^{3,4}

¹ *Purple Mountain Observatory, Chinese Academy of Sciences, Nanjing, 210008, China*

² *National Astronomical Observatories, Chinese Academy of Sciences, China*

³ *Department of Astronomy, Nanjing University, Nanjing, 210093, China*

⁴ *LCRHEA, IHEP, CAS, Beijing, China*

1 February 2008

All correspondence please send to:

D.M. Wei
Purple Mountain Observatory
Chinese Academy of Sciences
Nanjing, 210008
P.R. China

email: dmwei@pmo.ac.cn

fax: 8625-3307381

arXiv:astro-ph/0012007v2 25 Feb 2002

ABSTRACT

Whether gamma-ray bursts are highly beamed or not is a very important question, since it has been pointed out that the beaming will lead to a sharp break in the afterglow light curves during the ultra-relativistic phase, with the breaking point determined by $\Gamma \sim 1/\theta_0$, where Γ is the bulk Lorentz factor and θ_0 is the initial half opening angle of the ejecta, and such a break is claimed to be present in the light curves of some GRBs. In this paper we will examine whether all the observed breaks in GRB afterglow light curves can be explained by jet effects. Here we present a detailed calculation of the jet evolution and emission, and have obtained a simple formula of bulk Lorentz factor evolution. We show that the light curves are very smoothly steepened by jet effect, and the shape of the light curve is determined by only one parameter — $(E/n)^{1/8}\theta_0^{3/4}$, where E and n are the fireball energy and surrounding medium density. We find that for GRB990123 and GRB991216, the jet model can approximately fit their light curves, and the values of $(E_{51}/n_1)^{1/8}\theta_0^{3/4}$ are about 0.17 and 0.22 respectively. While for GRB990510, GRB000301c, GRB000926 and GRB010222, their light curves cannot be fitted by the jet model, which suggests that the breaks may be caused by some other reasons, jet should be not the unique reason.

Key words: gamma rays: bursts — shock waves — ISM: jets and outflows

1 INTRODUCTION

Multiwavelength follow-up of gamma-ray burst afterglows has revolutionized GRB astronomy in recent years, yielding a wealth of information about the nature of GRBs (Klose 2000; Castro-Tirado et al. 1999 and references therein). The afterglows can well be explained as the emission from a relativistic blast wave which decelerates when sweeping up interstellar medium. The dynamical evolution of GRB fireballs and the emission features have been studied by many authors (e.g. Sari 1997; Meszaros, Rees & Wijers 1998; Wei & Lu 1998a, 2001a; Huang et al. 2000a; Sari, Piran & Narayan 1998; Wijers, Rees & Meszaros 1997).

One of the most important questions to the nature of GRBs is the total energy released in the event, which depends on two factors: the distance scale and the opening angle of GRB ejecta. The distance scale of several GRBs have been derived through measuring the redshift in optical observations, while the question whether the gamma-ray emission is isotropic or not remains uncertain. The observations show that, if the gamma ray emission is isotropic, then for some GRBs their total energy will be too large, for example, the isotropic energies of GRB971214 and GRB990123 are 3×10^{53} ergs and 3.4×10^{54} ergs respectively (Kulkarni et al. 1998, 1999). Such a crisis encountered by this extreme large energy forced some people to think that the GRB emission must be highly collimated in order to reduce the total energy.

Then how can we tell a jet from an isotropic fireball? Rhoads (1997, 1999) has pointed out that the lateral expansion of the relativistic jet will produce a sharp break in the afterglow light curves, and such breaks are also claimed to be present in the light curves of GRB990123 (Kulkarni et al. 1999; Castro-Tirado et al. 1999), GRB 990510 (Harrison et al. 1999; Stanek et al. 1999), GRB000301c (Rhoads & Fruchter 2001; Masetti et al. 2000) and GRB000926 (Sagar et al. 2001; Price et al. 2001), GRB991216 (Halpern et al. 2000), GRB010222 (Masetti et al. 2001; Stanek et al. 2001). Here we give a detailed analysis of the jet evolution and emission under the relativistic case to examine whether the jet model can account for all the observed breaks in GRB afterglows, and find that for four bursts, GRB990510, GRB010222, GRB000926 and GRB000301C, their afterglow light curves cannot be explained by jet effects. In next section we discuss the jet evolution and emission in the relativistic regime, in section 3 we fit six GRBs' afterglow light curves, and finally we give some discussions and conclusions.

2 THE EVOLUTION AND EMISSION OF JET

Here we follow our previous paper (Wei & Lu 2000b) to calculate the emission flux from the jet. We assume that the radiation is isotropic in the comoving frame of the ejecta, the radiation cone is uniquely defined by the angular spherical coordinates (θ, ϕ) , here θ is the angle between the line of sight (along z -axis) and the symmetry axis, and ϕ is the azimuthal angle. Because of cylindrical symmetry, we can assume that the symmetry axis of the cone is in the $y - z$ plane. In order to see more clearly, let us establish an auxiliary coordinate system (x', y', z') with the z' -axis along the symmetry axis of the cone and the x' parallel the x -axis. Then the position within the cone is specified by its angular spherical coordinates θ' and ϕ' ($0 \leq \theta' \leq \theta_j$, $0 \leq \phi' \leq 2\pi$, θ_j is the jet opening half angle, which increases with time). It can be shown that the angle Θ between a direction (θ', ϕ') within the cone, and the line of sight satisfies $\cos\Theta = \cos\theta\cos\theta' - \sin\theta\sin\theta'\sin\phi'$. Then the observed flux is

$$F_\nu = \int_0^{2\pi} d\phi' \int_0^{\theta_j} \sin\theta' d\theta' D^3 I'(\nu D^{-1}) \frac{r^2}{d^2} \quad (1)$$

where $D = [\Gamma(1 - \beta\cos\Theta)]^{-1}$ is the Doppler factor, $\beta = (1 - \Gamma^{-2})^{1/2}$, $\nu = D\nu'$, $I'(\nu')$ is the specific intensity of synchrotron radiation at ν' , and d is the distance of the burst source. Here the quantities with prime are measured in the comoving frame. For simplicity we have ignored the relative time delay of radiation from different parts of the cone.

It is widely believed that the electrons have been accelerated by the shock to a power law distribution $n_e(\gamma) \propto \gamma^{-p}$ for $\gamma_{min} \leq \gamma \leq \gamma_{max}$, and consider the synchrotron radiation of these electrons, we can obtain the observed flux

$$F_\nu \propto f(\Gamma) z T^3 n^{(p+5)/4} \xi_B^{(p+1)/4} \nu^{-(p-1)/2} \quad (2)$$

where T is the time measured in the observer frame, $f(\Gamma) = \Gamma^{-(p+7)/2} (\Gamma-1)^{(p+1)/4} y(\Gamma)^{(p+5)/4} \gamma_{min}^{p-1} \beta^3$, $y(\Gamma) = \frac{\hat{\gamma}\Gamma+1}{\hat{\gamma}-1}$, $\hat{\gamma}$ is just the ratio of specific heats, $\gamma_{min} = \xi_e(\Gamma-1) \frac{m_p}{m_e} \frac{p-2}{p-1}$, $m_p(m_e)$ is the mass of proton (electron), and when the line of sight is along with the jet symmetry axis, $z = \beta^{-1}[(1-\beta)^{-(p+9)/2} - (1-\beta\mu_j)^{-(p+9)/2}]$, $\mu_j = \cos\theta_j$, n is the surrounding medium density, ξ_B is the energy fraction occupied by magnetic field. In the relativistic case, we have

$$F_\nu \propto \Gamma^{2p+6} T^3 [1 - (\frac{1}{1 + \Gamma^2 \theta_j^2})^{(p+9)/2}] n^{(p+5)/4} \xi_B^{(p+1)/4} \quad (3)$$

Now let us discuss the jet evolution. For an adiabatic relativistic jet expanding in surrounding medium, the evolution equation of energy conservation is

$$\Gamma^2 N m_p c^2 = E \quad (4)$$




Figure 1. The evolution of $\Gamma\theta_0$ with T_* . The circles are the numerical results, and the solid line is our fit results (eq.(6)).

For constant density, the total particle numbers $N = nV = 2\pi nr^3(1 - \cos\theta_j)/3 \simeq \pi nr^3\theta_j^2/3$ for $\theta_j \ll 1$. In the relativistic case, $r \simeq 4\Gamma^2 cT$, so we have

$$\Gamma = 372 \left(\frac{E_{51}}{n_1} \right)^{1/8} \theta_j^{-1/4} T^{-3/8} \quad (5)$$

where E_{51} is the fireball energy in units of 10^{51} ergs, n_1 is the surrounding medium density in units of 1 atom cm^{-3} . The jet half opening angle $\theta_j = \theta_0 + \theta' \simeq \theta_0 + \frac{2}{5} \frac{c_s}{c} \frac{1}{\Gamma}$, where c_s is the expanding velocity of ejecta material in its comoving frame, and for relativistic expanding material it is appropriate to take c_s to be the sound speed $c_s = c/3^{1/2}$ (Rhoads 1997, 1999). Then we can obtain the evolution of jet Lorentz factor $\Gamma\theta_0(1 + \frac{2}{5} \frac{c_s}{c} \frac{1}{\Gamma\theta_0})^{1/4} = 372 \left(\frac{E_{51}}{n_1} \right)^{1/8} \theta_0^{3/4} T^{-3/8}$. So we see that, if the value of $(\frac{E_{51}}{n_1})^{1/8} \theta_0^{3/4}$ is given, then we can calculate the variation of $\Gamma\theta_0$ with time T , and here we obtained a simple fitted formula

$$\Gamma\theta_0 = 5.2 T_*^{-\alpha(T_*)} \quad (6)$$

where $T_* = (\frac{E_{51}}{n_1})^{-1/3} \theta_0^{-2} T_{day}$, T_{day} is the observed time in units of 1 day, and $\alpha(T_*) = 0.375 + 0.075 \frac{(T_*/80000)^{0.29}}{1 + (T_*/80000)^{0.29}}$. It should be noted that the index α is a function of T_* , so the relation between $\Gamma\theta_0$ and T_* is not simply determined by α , the actual decay index k should be calculated as $k = d \ln(\Gamma\theta_0) / d \ln T_* = -(\alpha + \ln T_* d\alpha / d \ln T_*)$. Fig.1 gives the numerical results of $\Gamma\theta_0$ evolution and our fitted results. Equation (6) can be easily used to evaluate the evolution of bulk Lorentz factor, and we can derive the value of $(E/n)^{1/8} \theta_0^{3/4}$ through fitting the afterglow light curve.




Figure 2. The R-band light curve of the GRB990123 afterglow, the contribution of host galaxy has been subtracted. The solid line represents our best fit. The fit parameters are: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.17 \pm 0.08$, $p = 2.49 \pm 0.25$, the reduced $\chi^2 = 2.2$.

3 FITTING RESULTS

Based on the results described above, now we fit six GRBs' afterglow light curves in which the breaks are present, they are GRB990123, GRB990510, GRB000301c, GRB000926, GRB991216 and GRB010222, the results are as follows.

3.1 GRB990123

GRB990123 was the brightest GRB seen by BeppoSAX to date, it is in the top 0.3% of all bursts if ranked by the observed fluence, its redshift is estimated to be $z = 1.6$, corresponding to a luminosity distance $d_L \simeq 12$ Gpc, and the isotropic γ -ray energy $E_\gamma \approx 3.4 \times 10^{54}$ ergs (Kulkarni et al. 1999). The observed spectral slope between the optical band and the X-ray wavelengths was $\beta_{OX} = -0.68 \pm 0.05$, corresponding to the electron distribution index p between 2.3 and 2.5 (Castro-Tirado et al. 1999). Fig.2 illustrates our best fit to the R band light curve, the contribution from the host galaxy has been subtracted. The best fit parameters are: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.17 \pm 0.08$, $p = 2.49 \pm 0.25$, and the reduced $\chi^2 = 2.2$. We see that jet expansion can explain the observed light curve.

3.2 GRB000926

GRB000926 was a burst lasted about 25 s, its redshift is about $z = 2.066$, yielding the luminosity distance $d_L = 16.9$ Gpc, and the isotropic energy $E_\gamma \simeq 2.5 \times 10^{53}$ ergs. The value of spectral index in the X-ray –optical region is about $\beta = -0.8$ or -0.9 , corresponding to the electron distribution index p between 2.6 and 2.8 (Sagar et al. 2001). Fig.3 gives our best




Figure 3. The R-band light curve of the GRB000926 afterglow, the contribution of host galaxy has been subtracted. The solid line represents our best fit. The fit parameters are: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.087 \pm 0.053$, $p = 2.61 \pm 0.36$, and the reduced $\chi^2 = 6$.

fit to the R band light curve, the contribution from the host galaxy has been subtracted. The fitted parameters are: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.087 \pm 0.053$, $p = 2.61 \pm 0.36$, and the reduced $\chi^2 = 6$, so we see that the jet model cannot fit the light curve well.

3.3 GRB991216

GRB991216 was one of the brightest γ -ray bursts detected by BATSE, with a fluence of 2.56×10^{-4} ergs cm^{-2} above 20.6 KeV (Kippen 1999). Its duration is about 15 seconds, and the redshift is about $z = 1.02$, yielding the isotropic gamma-ray energy about 6.7×10^{53} ergs. The observed spectral slope between the optical band and the X-ray wavelengths was $\beta_{OX} = -0.74 \pm 0.05$, corresponding to the electron distribution index p between 2.4 and 2.6 (Halpern et al. 2000). Fig.4 illustrates our best fit to the R band light curve, the contribution from the host galaxy has been subtracted. The best fit parameters are: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.22 \pm 0.04$, $p = 2.51 \pm 0.08$, and the reduced $\chi^2 = 1.5$. We see that jet expansion can explain the observed light curve well.

3.4 GRB990510

GRB990510 was also a strong burst with duration of about 100 s, its redshift is $z = 1.619 \pm 0.002$, corresponding to the luminosity distance $d_L \simeq 12$ Gpc, and the isotropic energy $E_\gamma \approx 2.9 \times 10^{53}$ ergs (Vreeswijk et al. 1999). The observed optical spectral index was $\beta = -0.61 \pm 0.12$, corresponding to the electron distribution index p between 2.1 and 2.3 (Stanek et al. 1999). However, we find this burst cannot be fitted by simple jet model, since the




Figure 4. The R-band light curve of the GRB991216 afterglow, the contribution of host galaxy has been subtracted. The solid line represents our best fit. The fit parameters are: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.22 \pm 0.04$, $p = 2.51 \pm 0.08$, and the reduced $\chi^2 = 1.5$.




Figure 5. The R-band light curve of the GRB990510 afterglow. The solid line represents our best fit. The fit parameters are: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.096 \pm 0.004$, $p = 1.87 \pm 0.03$, $t_c = 438 \pm 25$ days, the reduced $\chi^2 = 7.8$.

reduced $\chi^2 = 23.5$. One possible way to improve the fitting results is that the cooling frequency ν_c crosses the observed band around break time. Similar to eq.(3), we can obtain the observed flux for $\nu > \nu_c$ in the relativistic case

$$F_\nu \propto \Gamma^{2p+4} T^2 \left[1 - \left(\frac{1}{1 + \Gamma^2 \theta_j^2} \right)^{(p+8)/2} \right] \quad (7)$$

Using eqs.(3)(7) and $\Gamma\theta_0$ evolution, we re-fit the afterglow light curve of GRB990510, the parameters are: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.096 \pm 0.004$, $p = 1.87 \pm 0.03$, $t_c = 438 \pm 25$ days (where t_c is the time $T_\star$ when the cooling frequency ν_c crosses the observed band), the reduced $\chi^2 = 7.8$. From Fig.5 we see that the jet model still cannot explain the light curve well.




Figure 6. The R-band light curve of the GRB010222 afterglow. The solid line represents our best fit. The fit parameters are: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.169 \pm 0.018$, $p = 2.01 \pm 0.04$, $t_c = 105 \pm 35$ days, the reduced $\chi^2 = 5.1$.

3.5 GRB010222

GRB010222 was detected simultaneously by the Gamma-Ray Burst Monitor (GRBM) and the Wide Field Camera (WFC) instruments on BeppoSAX, the 40 – 700 KeV fluence was as bright as $(9.3 \pm 0.3) \times 10^{-5}$ ergs cm^{-2} , only be surpassed by GRB990123. The redshift of this burst was determined at $z = 1.477$ (Jha et al. 2001). The spectral index of optical data is $\beta = -0.88 \pm 0.10$, corresponding to the electron distribution index p between 2.5 and 3 (Stanek et al. 2001). However, this burst also cannot be fitted by simple jet model, since the reduced $\chi^2 = 15.7$. If we consider the case that the cooling frequency ν_c crosses the observed band around break time, we obtain the following fit parameters: $(E_{51}/n_1)^{1/8}\theta_0^{3/4} = 0.169 \pm 0.018$, $p = 2.01 \pm 0.04$, $t_c = 105 \pm 35$ days, the reduced $\chi^2 = 5.1$. From Fig.6 we see that the jet model can approximately fit the light curve. However, it should be noted that the fit parameter p is obviously smaller than that from spectra measure, and in addition, Masetti et al. (2001) has pointed out that the constancy of the optical spectral shape excluded that the ν_c has crossed the optical band during the observations, so we think that the jet model still cannot explain the light curve.

3.6 GRB000301c

GRB000301c was a peculiar burst with duration of about 10 s, there are short term flux variability in its optical light curve, and besides this, the overall light curve shows sharp break (Sagar et al. 2000; Masetti et al. 2000). Its redshift is $z = 2.0335 \pm 0.0003$, corresponding to the luminosity distance $d_L = 16.6$ Gpc, and the isotropic energy $E_\gamma \simeq 3.4 \times 10^{53}$ ergs. The

spectral index is about $\beta = -0.8$ or -0.9 , corresponding to the electron distribution index p between 2.6 and 2.8 (Sagar et al. 2000). However, this burst cannot be fitted by simple jet model, since the reduced $\chi^2 = 31$. If we consider the cooling frequency ν_c crosses the observed band around break time, the reduced χ^2 value is still very high, $\chi^2 = 25$, so this burst also cannot be explained by jet model.

4 DISCUSSION AND CONCLUSION

Whether gamma ray bursts are beamed or not is a very important question, since it is related with the energy sources of GRBs, and the afterglow observations provide a very good chance to study this question. Rhoads (1997, 1999) pointed out that the sideways expansion of the relativistic jet would cause the blast wave to decelerate more quickly, leading to a sharp break in the afterglow light curve. However, some numerical calculations show that such break is much weaker and smoother than the prediction (Moderski et al. 2000; Panaitescu & Mészáros 1999; Kumar & Panaitescu 2000). We have also pointed out that the steepening of the afterglow light curves can be observed only when the beaming angle is very small, $\theta_0 < 0.1$ (Wei & Lu 2000b), but in that paper we fixed the Lorentz factor $\Gamma_0 = 300$.

In this paper we reanalyse the jet evolution and emission in detail, and we find that the effect of jet expansion on the afterglow light curve can be described by only one parameter — $(E/n)^{1/8}\theta_0^{3/4}$. From equations (3) and (7) it is obvious that, when $\Gamma\theta_j \gg 1$, the observed flux $F_\nu \propto T^{-3(p-1)/4}$ for $\nu < \nu_c$ or $F_\nu \propto T^{-3(p-2)/4}$ for $\nu > \nu_c$, while when $\Gamma\theta_j \ll 1$, the flux $F_\nu \propto T^{-3p/4}$ for $\nu < \nu_c$ or $F_\nu \propto T^{-(3p+1)/4}$ for $\nu > \nu_c$ if no lateral expansion, and $F_\nu \propto T^{-p}$ if lateral expansion is important, these are the same as the previous analytical results, which claim that there will be two breaks in the afterglow light curve. However, from equations (3) and (7) we see that the afterglow light curve is steepened gradually and smoothly, which depends on the variation of $\Gamma\theta_j$.

The breaks predicted by theoretical models have been observed in some GRBs' afterglow light curves, and have been generally considered as evidence for collimation of the relativistic GRB ejecta. We have fitted six GRBs' afterglow light curves in which the breaks are present, and find that, for GRB990123 and GRB991216, their light curves can be approximated fitted by the jet model, but for GRB990510, GRB000301c, GRB000926 and GRB010222, their light curves cannot be fitted by the jet model. So we conclude that, although the lateral expansion of the relativistic jet can lead to a break in the afterglow light curve, this may be not the

unique reason, there should be some other reasons to cause the break. For example, the transition from relativistic phase to non-relativistic phase of the blast wave may cause such break (Dai & Lu 1999; Huang et al. 2000b); the effects of inverse Compton scattering can flatten or steepen the light curves (Wei & Lu 1998b, 2000a); the curved emission spectra (not a simple power-law as usually assumed) can also produce a break in the light curves (Wei & Lu 2002).

From fitting we have obtained the values of $(E_{51}/n_1)^{1/8}\theta_0^{3/4}$. As an approximation, this value can also be regarded as the value of $\theta_0^{3/4}$, since it is very weakly dependent on the value of (E_{51}/n_1) . Therefore from the fitting we can estimate the opening half angle of GRBs, for GRB990123 $\theta_0 \approx 0.094$, and for GRB991216 $\theta_0 \approx 0.13$. In addition, from equation (5) we see that, if the value of $(E_{51}/n_1)^{1/8}\theta_0^{3/4}$ is known, then we can also get the value of $\Gamma_0\theta_0 t_0^{3/8}$, here t_0 is the deceleration time. In a few bursts, soft X-ray emission has been observed from the end of the GRB phase, indicating that the external shock had already set in by the end of the GRB (at t_γ) (Giblin et al. 1999). In other cases there is no detectable X-ray emission after the GRB, suggesting that $t_\gamma < t_0$ (Pian et al. 2001). In order to constrain $\Gamma_0\theta_0$, we assume that the observed GRB duration is a good measure of t_0 (see also Panaitescu & Kumar 2001), and equation (5) shows that Γ_0 has a moderate dependence on t_0 , thus the error due to this assumption is likely not too large. Furthermore, using the values of θ_0 derived above, we can give an estimate of the initial Lorentz factor, we find for GRB990123 $\Gamma_0 \approx 130$, and for GRB991216 $\Gamma_0 \approx 228$. These values of Γ_0 seem reasonable since it is widely believed that the initial fireball Lorentz factor should be larger than 100 in order to avoid photon-photon annihilation.

Here we only consider the case that the line of sight is along the jet axis, since only this can give an analytical results. For the case that the line of sight is off the jet axis but still within the cone, we have calculated the light curves using equation (1), and found that the overall shape of the light curve is similar to the case $\theta = 0$, but the break is more smooth, as suggested by Ghisellini & Lazzati (2000), who did not consider the lateral expansion.

In summary, our conclusion is that, not all the breaks in the afterglow light curves can be explained by jet model, so the steepening of the light curves may be caused by varied reasons, jet should be not the unique reason. But if the light curve can be fitted by jet model, then we can obtain the value of $(E_{51}/n_1)^{1/8}\theta_0^{3/4}$, and furthermore we can give a constraint on the values of Γ_0 and θ_0 .

5 ACKNOWLEDGEMENTS

We would like to thank the referee for several important comments that improved this paper. Also thanks to Drs. Z.G.Dai and Y.F. Huang for helpful discussions. This work is supported by the National Natural Science Foundation (10073022 and 19973003) and the National 973 Project on Fundamental Researches of China (NKBRF G19990754).

REFERENCES

- Castro-Tirado, A.J., et al., 1999, *Science*, 283, 2069
- Dai, Z.G., Lu, T., 1999, *ApJ*, 519, L155
- Giblin, T., et al., 1999, *ApJ*, 524, L47
- Ghisellini, G., Lazzati, D., 2000, *MNRAS*, 309, L7
- Halpern, J.P., et al., 2000, *ApJ*, 543, 697
- Harrison, F.A., et al., 1999, *ApJ*, 523, 121
- Huang, Y.F., et al., 2000a, *ApJ*, 543, 90
- Huang, Y.F., Dai, Z.G., Lu, T., 2000b, *A&A*, 355, L43
- Jha, S., et al., 2001, *ApJ*, 554, L155
- Kippen, R.M., 1999, *GCN Circ.* 504
- Klose, S., 2000, *Rev. Modern Astron.*, 13, 129
- Kulkarni, S.R., et al., 1998, *Nature*, 393, 35
- Kulkarni, S.R., et al., 1999, *Nature*, 398, 389
- Kumar, P., Panaitescu, A., 2000, *ApJ*, 541, L9
- Masetti, N., et al., 2000, *A&A*, 359, L23
- Masetti, N., et al., 2001, *A&A*, 374, 382
- Mészáros, P., Rees, M.J., Wijers, R., 1998, *ApJ*, 499, 301
- Moderski, R., Sikora, M., Bulik, T., 2000, *ApJ*, 529, 151
- Panaitescu, A., Mészáros, P., 1999, *ApJ*, 526, 707
- Panaitescu, A., Kumar, P., 2001, submitted to *ApJ*, astro-ph/0109124
- Pian, E., et al., 2001, *A&A*, 372, 456
- Price, P.A., et al., 2001, *ApJ*, 549, L7
- Rhoads, J.E., 1997, *ApJ*, 487, L1
- Rhoads, J.E., 1999, *ApJ*, 525, 737
- Rhoads, J.E., Fruchter, A.S., 2001, *ApJ*, 546, 117
- Sagar, R., et al., 2000, *Bull. Astron. Soc. India*, 28, 499 (astro-ph/0004223)
- Sagar, R., et al., 2001, *Bull. Astron. Soc. India*, 29, 1 (astro-ph/0010212)
- Sari, R., 1997, *ApJ*, 494, L49
- Sari, R., Piran, T., Narayan, R., 1998, *ApJ*, 497, L17
- Stanek, K.Z., et al., 1999, *ApJ*, 522, L39
- Stanek, K.Z., et al., 2001, *ApJ*, in press (astro-ph/0104329)
- Vreeswijk, P.M., et al., 1999, *GCN* 310
- Wei, D.M., Lu, T., 1998a, *ApJ*, 499, 754
- Wei, D.M., Lu, T., 1998b, *ApJ*, 505, 252
- Wei, D.M., Lu, T., 2000a, *A&A*, 360, L13

Wei, D.M., Lu, T., 2000b, ApJ, 541, 203

Wei, D.M., Lu, T., 2001a, MNRAS, 320, 37

Wei, D.M., Lu, T., 2002, A&A, 381, 731 (astro-ph/0107371)

Wijers, R.A.M.J., Rees, M.J., Mészáros, P., 1997, MNRAS, 288, L51